



1. *Cylindrella*. Ghiesbreghtii Pf^r 2. *C. turris* Pf^r 3. *C. clava* Pf^r
4. 5. *Helix* muta Pf^r, and var. 6. 7. *H. Lombaei* Pf^r, and var.
8. *H. Grevillei* Pf^r

tura obliqua, subangulato-circulari; perist. subduplice interno vix interrupto, adnato, externo patente, concentrica striatulo, latere sinistro dilatato, fornicatim reflexo.—Operc. planum, fulvum.

Diam. maj. 12, min. 10, alt. 10 mill.

17. **HELICINA GRATIOSA**, Pfr. *H. testa subgloboso-turbinata, tenuiuscula, striatula, striolis spiralibus obsoletissime notata, nitida, pellucida, fuscescenti-rubra vel lutea; spira conoidea, acutiuscula; anfr. 5 convexiusculis, ultimo subcarinato, antice rotundato; columella subrecedente, alba, callum emittente albidum; apertura obliqua, subtriangulato-semicirculari; perist. undique breviter expanso.—Operc. tenue, solidum, carneum.*

Diam. maj. 9, min. $7\frac{1}{2}$, alt. fere 6 mill.

18. **HELICINA SUAVIS**, Pfr. *H. testa turbinata, tenui, nitidisima, superne striatula striisque spiralibus conferte notata, rubella, albido-marmorata, vel lutea aut carnea, pallide rubro variegata; spira conoidea, acuta; anfr. fere 5 convexiusculis, ultimo compresse et acute carinato, basi convexiore; columella brevi, basi subdentata, callum evittente crassiusculum diffusum; apertura obliqua, subtriangulari; perist. albo, margine supero breviter expanso, basali reflexiusculo.—Operc.?*

Diam. maj. $10\frac{1}{2}$, min. $8\frac{3}{4}$, alt. $6\frac{2}{3}$ mill.

Besides the described new species, there were in the same collection from the Admiralty Islands, *Helix Novæ Hibernicæ*, Quoy, *fringilla*, Pfr. (beautiful and large varieties), *migratoria*, Pfr., *Sachaleensis*, Pfr., *motacilla*, Pfr., *Cleryi*, Recl., *helicinaoides*, Jacq., *pyxis*, Hinds, *Grimardi*, Desh., *breripila*, Pfr., *Pfeifferi*, Phil., &c.

5. DESCRIPTIONS OF THIRTY-THREE NEW SPECIES OF LAND-SHELLS, FROM THE COLLECTION OF H. CUMING, ESQ.
BY DR. L. PFEIFFER.

(Mollusca, Pl. XXXVI.)

1. **HELIX GLORIOSA**, Pfr. *H. testa imperforata, depresso ovata, crassa, ponderosa, oblique conferte plicato-striata lirisque crebris obtusis subreguluribus cincta, fulvida, strigis singulis pallidis et saturioribus notata; spira convexa, obtusa, nuda, sub lente granulata, apice subtilissimo; anfr. $3\frac{1}{2}$ rapide accrescentibus, ultimo perinflato; apertura diagonali, lunato-ovali, intus rubella; perist. crassissimo, breviter expanso, marginibus callo crasso junctis, columellari adnato.*

Diam. maj. 64, min. 50, alt. 39 mill.

Hab. Madagascar.

2. **HELIX DUCTILIS**, Pfr. *H. testa umbilicata, discoidea, tenui,*
No. CCCXXV.—PROCEEDINGS OF THE ZOOLOGICAL SOCIETY.

sub lente confertim striatula, albido-hyalina; spira plana; anfr. fere 5 subplanis, sensim accrescentibus, ultimo non descendente, depresso, peripheria rotundato, basi vix converxiore; umbilico perspectivo, $\frac{1}{4}$ diametri superante; apertura obliqua, late lunari; perist. simplice, recto, marginibus vix convergentibus, columellari brevi, verticali, cum basali subangulatim juncto.

Diam. maj. $6\frac{1}{2}$, min. $5\frac{1}{2}$, alt. 2 mill.

Hab. Drayton Range, North Australia (Mr. Stutchbury).

3. **HELIX STUTCHBURYI**, Pfr. *H. testa sub obtecte perforata, turbinato-globosa, tenui, superne sub lente minute granulata, pallide fulva, ad suturam et supra medium rufo-fasciata; spira convexo-conoidea, obtusula; anfr. 5 convexiusculis, ultimo rotundato, antice vix descendente, basi lævigato; apertura diagonali, rotundato-lunari; perist. tenui, marginibus subconvergentibus, dextro vix expansiusculo, basali breviter reflexo, columellari fornicatim dilatato, umbilicum angustissimum fere tegente.*

Diam. maj. $15\frac{1}{2}$, min. $13\frac{1}{2}$, alt. 10 mill.

Hab. Drayton Range, North Australia (Mr. Stutchbury).

4. **HELIX DELTA**, Pfr. *H. testa sub obtecte perforata, trochiformi, tenuiuscula, striatula et foveolata, opaca, carneo-albida; spira regulariter conica, apice acuta; anfr. 6 planis, ultimo non descendente, subacute carinato, basi convexiusculo; apertura fere diagonali, angulato-lunari; perist. simplice, margine dextro recto, basali reflexiusculo, columellari supra perforationem dilatato, patente.*

Diam. maj. 8, min. $7\frac{1}{2}$, alt. $6\frac{1}{2}$ mill.

Hab. Drayton Range, North Australia (Mr. Stutchbury).

5. **HELIX PLICULOSA**, Pfr. *H. testa umbilicata, turbinato-globulosa, tenuiuscula, superne confertim plicosa, diaphana, parum nitida, rubello-cornea; spira conoidea, vertice subtili; anfr. $4\frac{1}{2}$ convexiusculis, ultimo supra medium subangulato, antice rotundato, vix descendente, basi inflato, circa umbilicum angustum, pervium subcompresso; apertura obliqua, lunato-rotundata; perist. albido, marginibus subconniventibus, dextro expansiusculo, columellari sursum dilatato, cum basali angulum obtusum formante.*

Diam. maj. 16, min. 13, alt. 10 mill.

Hab. Drayton Range, North Australia (Mr. Stutchbury).

6. **HELIX CASTRENSIS**, Pfr. *H. testa imperforata, conoideo-semiglobosa, solida, rugoso-plicatula et irregulariter subgranulata, pallide fulvida, strigis fuscis crebris regulariter radiata; spira convexiusculo-conoidea, obtusula; anfr. 4 vix convexiusculis, ultimo medio subacute carinato, antice breviter descendente, inflato, basi pallidiore; apertura perobliqua, subtriangulari-lunari; perist. albo, margine dextro arcuato, breviter reflexo, basali dilatato, adnato, introrsum dentibus 2 obtusis plicæformibus munito.*

Diam. maj. 25, min. 21, alt. 15 mill.

Hab. West Indies.

7. **HELIX LINDSTEDTI**, Pfr. *H. testa sinistrorsa, perforata, conoideo-leuiculari, carinata, tenui, diaphana, superne undulato-striata et minutissime granulata, parum nitida, rubello-lutescente; spira regulariter elevata, obtusa; sutura linearis; anfr. 6 planiusculis, sensim accrescentibus, ultimo non descendente, supra medium acute carinato, basi convexo, nitidissimo, non decussato, circa perforationem non apertam albido; apertura fere diagonalis, securiformis; perist. simplice, recto, margine columellari in laminam brevem revolutam dilatato.*

Diam. maj. $35\frac{1}{2}$, min. 31, alt. 14 mill.

Hab. Malacca (Rev. F. W. Lindstedt).

8. **HELIX BOURGUIGNATI**, Pfr. *H. testa umbilicata, globoso-depressa, tenui, minutissime malleato-striata, parum nitente, diaphana, carneus; spira breviter conoidea, vertice subtili, luteo; anfr. 6 convexiusculis, regulariter accrescentibus, ultimo rotundato, antice vix descendente; umbilico angusto, vix pervio; apertura obliqua, rotundato-lunari; perist. intus albolabiato, marginibus vix convergentibus, dextro recto, basali reflexiusculo, ad umbilicum dilatato, patulo.*

Diam. maj. 19, min. 16, alt. 11 mill.

Hab. Crimea.

9. **HELIX ADELAIDÆ**, Pfr. *H. testa umbilicata, depressa, carinata, tenuiuscula, striata, alba, fascia fuscula supera ornata; spira vix elevata, vertice obtusiusculo, corneo; anfr. 4 convexiusculis, ultimo non descendente, supra peripheriam subacute carinato, basi inflato; umbilico perspectivo, $\frac{1}{4}$ diametri fere æquante; apertura fere diagonalis, lunato-rotundata, ad carinam vix angulata; perist. recto, intus levissime labiato, marginibus conniventibus, columellari subpatulo.*

Diam. maj. 6, min. 5, alt. fere 3 mill.

Hab. Adelaide, Australia.

10. **HELIX CHIONODISCUS**, Pfr. *H. testa subobtecte umbilicata, conoideo-lentiformi, carinata, solida, rugoso-striatula, opaca, nivea; spira convexiusculo-conoidea, nucleo læviusculo, obtuso; anfr. 5 vix convexiusculis, ultimo antice breviter deflexo, infra suturam linearem turgidulo, peripheria magis minusve acute carinato, basi convexo; apertura diagonalis, rhomboe-lunari; perist. subcalloso, marginibus subparallelis, dextro recto, basali perarcuato, reflexo, versus umbilicum sensim dilatato, lamina lata adnata illum fere claudente.*

Diam. maj. 26, min. 23, alt. 13 mill.

Hab. Crimea.

11. **HELIX GREVILLEI**, Pfr. (Pl. XXXVI. fig. 8.) *H. testa imperforata, conica, solida, striatula et plicis validis oblique antorsum descendenteribus, interdum malleato-interruptis munita, castanea, strigis sparsis nigricantibus notata; spira conoidea,*

apice rosea, acutiuscula; anfr. 4½ convexiusculis, rapide accrescentibus, penultimo subtus albo-callosa, ultimo antice descendente, peripheria obtuse sed distincte angulata; apertura pero obliqua, truncato-oblonga, intus pallide cæruleo-rosea; perist. nigro, expanso et reflexo, marginibus parallelis, callo nigro juncatis, columellari dilatato, plano, adnato, dextrorsum subdentato, ad axin albido.

Diam. maj. 49–58, min. 36–42, alt. 32–39 mill.

Hab. Ceylon (Mr. Thwaites).

12. ENNEA INSIGNIS, Pfr. *E. testa breviter rimata, ovato-oblonga, tenuissima, oblique confertim striata, striisque spiralibus versus basin evanescientibus decussatula, nitida, pellucida, cereo-hyalina; spira ovoidea, apice obtusa; sutura submarginata; anfr. 7 convexiusculis, penultimo supra aperturam subplanato, ultimo ¾ longitudinis formante, antice arcuatim ascendente, juxta rimam sub-compresso; columella recedente, dentato-plicata; perist. tenui, albo, undique expanso, murgine columellari dilatato, patente.*

Long. 37, diam. 20 mill.

Hab. Gaboon, Africa (Mr. Auboy).

13. BULIMUS PALAVANENSIS, Pfr. *B. testa imperforata, oblonga, solida, striatula, epidermide fusco-cinerea, custaneo varie strigata, sursum detrita munita; spira convexiusculo-conica, apice obtusula; anfr. 5–5½ modice convexis, ultimo spiram subæquante vel breviore; columella substricta, albida; apertura obliqua, truncato-ovali, intus griseo-cæruleo-rosea; perist. caloso, nigricante, breviter reflexo.*

Long. 48–49, diam. 27 mill.

Hab. Palawan (Dr. Traill).

14. BULIMUS LIBROSUS, Pfr. *B. testa imperforata, ovato-oblonga, solida, striatula, epidermide subhydrophana, libro-cinerea, fusco irregulariter strigata et ad suturam ocellata obducta; spira convexiusculo-conica, obtusa; anfr. 5½ planiusculis, ultimo spira vix breviore, epidermide decidua circa columellam pallidam, compres-sam, substrictum saturate castanea, nitida; apertura obliqua, ovali oblonga, intus cæruleo-rosea; perist. castaneo-nigro, subin-crassato, revoluto, margine dextro leviter arcuato.*

Long. 40, diam. 21 mill.

Hab. Palawan (Dr. Traill).

15. BULIMUS LINDSTEDTI, Pfr. *B. testa sinistrorsa, imperforata, ovato-conica, solida, striatula, nitida, candida; spira elongato-conica, apice obtusula; anfr. 6½ convexiusculis, ultimo ¾ longitudinis subæquante, obsolete angulata, antice rotundato; columella subverticali, vix torta; apertura fere diagonali, subsemi-circulari; perist. leviter incrassato, expanso, marginibus callo concolore juncatis.*

Long. 39, diam. 17 mill.

Hab. Malacca (Rev. F. W. Lindstedt).

16. **BULIMUS PARALLELUS**, Pfr. *B. testa compresse umbilicata, fusiformi, tenuiuscula, striatula, fulva, epidermide albida subreticulata; spira elongato-conica, obtusa; unfr. 7 modice convexis, ultimo spira paulo breviore, basi compresso; columella superne leviter torto-plicata; apertura verticali, oblonga, lateribus parallelis, basi effusa, utrinque fusco-strigata; perist. tenui, albo, expanso, margine dextro supra medium impresso, columellar i latiore, sulco arcuato ab anfractu contiguo separato.*

Long. 22, diam. 7 mill.

Hab. St. Catherine's, Brazil.

17. **BULIMUS CATHARINÆ**, Pfr. *B. testa breviter rimata, subperforata, fusiformi, solidula, undique leviter punctato-rugulosa, albida, punctis corneis substrigatim conspersa; spira ventroso-turrita, apice acutiuscula; anfr. 8½ convexiusculis, ultimo ¼ longitudinis vix attingente, basi crista valida, compressa et pone illam minore munito, antice interne et externe interrupte nigro-strigato; apertura obliqua, subseptemdentata; plica 1 lamellæformi in parte, secunda subquadrangulari ad columellanam, tertia obliqua in latere sinistro baseos effusæ, 3 subæqualibus et 1 minuta in margine dextro; perist. albo, undique modice expanso.*

Long. 23-25, diam. 7½-8 mill.

Hab. St. Catherine's, Brazil.

18. **BULIMUS SUGILLATUS**, Pfr. *B. testa anguste umbilicata, oblongo-turrita, tenui, irregulariter plicato-striata, cornea, strigis subpunctatis opacis albis, singulisque rufis notata; spira elongato-conica, acuta; sutura levissime crenulata; anfr. 9 convexis, ultimo ⅔ longitudinis subæquante, basi subcompresso; columella leviter et stricte recedente; apertura parum obliqua, oblonga; perist. simplice, margine dextro recto, columellar i sursum dilatato, forniciatim reflexo.*

Long. 24, diam. 9½ mill.

Hab. Bolivia.

19. **BULIMUS GAYI**, Pfr. *B. testa subobtecte umbilicata, conicovata, solidula, fusca, fasciis angustis albis varic ornata; spira conica, obtusa; anfr. 5 modice convexis, ultimo spiram paulo superante, basi rotundato; columella leviter arcuata; apertura parum obliqua, truncato-ovali, intus albida; perist. simplice, margine dextro recto, columellar i perdilatato, umbilicum angustum fere tegente.*

Long. 27, diam. 16 mill.

Hab. Bolivia.

20. **BULIMUS SABATIERI**, Pfr. *B. testa subperforata, ovato-oblonga, tenuiuscula, sublavigata, irregulariter striatula, strigis undulatis pallide lutescentibus et saturate brunneis alternantibus picta; spira conica, sursum pallidiore, apice obtuso; anfr. 6 parum convexis, ultimo spira paulo breviore, infra medium obsolete angulato; columella stricta, violaceo-fusca; apertura parum*

obliqua, anguste elliptica; perist. simplice, recto, margine columellari sursum dilatato, reflexo, subadnato.

Long. 22, diam. 12 mill.; ap. 11 mill. longa, $5\frac{1}{2}$ lata.

Hab. Banks of the "Fleuve blanc," China? (Mr. Sabatier).

21. **BULIMUS DUTAILLYI**, Pfr. *B. testa subperforata, oblongo-turrita, tenui, confertim plicatula, nitida, albida, fasciis 6-7 interruplis spadiceis ornata; spira elongata, apice acuta; anfr. 7 convexiusculis, ultimo $\frac{2}{5}$ longitudinis subæquante, basi attenuato; columella vix arcuata, subrecedente; apertura vix obliqua, elliptico-oblonga; perist. simplice, recto, margine columellari papyraceo, superne reflexo, subadnato.*

Long. 31, diam. 12 mill.; ap. $13\frac{1}{2}$ mill. longa, $6\frac{1}{2}$ lata.

Hab. Brazils (Mr. Dutailly).

22. **BULIMUS PLICULATUS**, Pfr. *B. testa umbilicata, ovato-conica, tenuiuscula, plicis levibus, subvariciformibus sculpta, haud nitente, pallide grisea, strigis angustissimis albis et fuscis irregulariter picta; spira conica, acuta; anfr. 7 convexiusculis, ultimo spira paulo breviore, basi circa umbilicum mediocrem, rotundum subcompresso; columella substricta; apertura vix obliqua, elliptico-oblonga; perist. simplice, recto, margine columellari a basi dilatato, subfornicatum reflexo.*

Long. 23, diam. $11\frac{1}{3}$ mill.

Hab. Bolivia.

23. **BULIMUS CLOUÉI**, Pfr. *B. testa perforata, ovato-turrita, solidula, striata et interdum submalleata, alba, strigis, maculis et punctis corneo-fascis irregulariter notata; spira elongato-conica, acutiuscula; anfr. 7 modice convexis, ultimo $\frac{3}{7}$ longitudinis vix æquante, subangulato, circa perforationem non perviam corneo-areolato; columella breviter recedente; apertura obliqua, sinuato-ovali, intus fuscula; perist. breviter expanso, margine columellari superne dilatato, reflexo, tum angulo obtuso ad basin descendente.*

Long. 22, diam. 10 mill.

Hab. Brazil (Mr. Cloué).

24. **BULIMUS PUNCTICULATUS**, Pfr. *B. testa profunde et compresse umbilicata, ovato-conica, tenuiuscula, sublævigata, nitida, albida, punctis pellucidis raris conspersa; spira elongato-conica, acutiuscula; anfr. 7 planiusculis, summis lutescentibus, ultimo spira breviore, antice ascendente, basi levissime compresso; columella subrecedente, leviter arcuata; apertura subverticali, oblongo-ovali; perist. simplice, margine dextro breviter expanso, columellari perdilatato, subflexuoso, patente.*

Long. 29, diam. $12\frac{1}{2}$ mill.

Hab. Bolivia.

25. **PARTULA ALABASTRINA**, Pfr. *P. testa compresse umbilicata, oblongo-conica, tenui, levissime striatula, vix nitidula, lutescenti-alabastrina; spira conica, obtusula; anfr. 5 $\frac{1}{2}$ convexis, ultimo*

spiram vix superante, basi attenuato, subcompresso; apertura parum obliqua, oblique truncato-oblonga; columella substricte recedente; perist. albo, marginibus callo tenuissimo junctis, dextro late expanso, columellari latissimo, patente.

Long. 23, diam. 11 mill.

Hab. Salomon's Islands.

26. CYCLOSTOMA (CYCLOTUS) LINDSTEDTI, Pfr. *C. testa umbilicata, depressa, subdiscoidea, distincte striata, strigis fulgurantibus nigro-fuscis et luteis, superne latis, subtus linearibus picta; spira vix elevata; anfr. 4 convexis, celeriter accrescentibus, ultimo terete, vix descendente; umbilico lato, $\frac{1}{3}$ diametri superante; apertura obliqua, subcirculari; perist. breviter adnato, duplicato; interno expansiusculo, externo albo, undique subæqualiter patente.—Operc. Cycloti.*

Diam. maj. 11, min. 9, alt. 4 mill.

Hab. Mount Ophir, Malacca (Rev. F. W. Lindstedt).

27. CATAULUS HÆMASTOMUS, Pfr. *C. testa breviter et profunde rimata, ovali-pyramidalis, tenuiuscula, conferte striatula, parum nitente, diaphana, citrina; spira convexiusculo-turrita, apice sub-acuta; sutura submarginata; anfr. 8 convexis, ultimo vix attenuato, basi axin excedente; carina umbilicali valida, compressa, antrorsum vix dilatata; periomphalo lato, costulato-striato; apertura subcirculari; perist. fusco-sanguineo, simplice, ad anfractum contiguum angustato, lateribus rectangule late patente et revoluto, basi subproducto, canali mediocri perforato.*

Long. 27-28, diam. 11-12 mill.

Hab. Ceylon (Mr. Thwaites).

28. HELICINA (ALCADIA) RHAMPHOSTYLA, Pfr. *H. testa conoideo-globosa, solida, sublævigata, albida vel lutescente; spira breviter conoidea, obtusula; anfr. 4 $\frac{1}{2}$, superis vix convexiusculis, ultimo magno, rotundato; columella lata, superne in cañum crassum, semicircularem dilatata, basi dextrorsum curvata; apertura obliqua, subsemicirculari; perist. expanso, tenui, intus calloso, margine dextro subrepando, basali sinu profundo ab extremitate rostriformi columellæ disjuncto, præterea plica pone columellam intrante munito.—Operc.?*

Diam. maj. 15 $\frac{1}{2}$, min. 13, alt. 12 mill.

Hab. —?

29. HELICINA NORFOLKENSIS, Pfr. *H. testa depresso-turbinata, solidula, leviter striata, albida; spira conoidea, versus apicem minute papillarem lutescente; anfr. 5 vix convexiusculis, ultimo antice rugoso, peripheria subangulata, basi plauisculo, callo granuloso nitido circumscripto et epidermide fulva decidua obducto; columella brevi, arcuata; apertura fere diagonali, subtriangulari-lunari; perist. simplice, recto, margine basali subincrassato.—Operc. testaceum, pallidum.*

Diam. maj. 14, min. 12, alt. 8 mill.

Hab. Norfolk Islands.

30. *HELICINA PICTELLA*, Pfr. *H. testa conoideo-depressa, tenui, sub lente tenuiter et subconferte lirata, parum nitida, diaphana, pallide cornea, ad suturam rubro-maculata; spira breviter conoidea; anfr. 4 convexiusculis, ultimo subangulato, basi callo tenui subcircumscripto obducto; columella brevissima, simplice, tenui; apertura parum obliqua, semiovali; perist. simplice, recto, margine basali angulum rectum cum columella formante.* —Operc. ?

Diam. maj. 4, min. $3\frac{1}{3}$, alt. 2 mill.

Hab. Norfolk Islands.

31. *HELICINA DRAYTONENSIS*, Pfr. *H. testa conoidea, solidula, ruguloso-striata striisque spiralibus nonnullis notata, parum nitida, cornea; spira conoidea, apice mucronulata, lutea; anfr. $4\frac{1}{2}$ planiusculis, ultimo subcarinato, busi convexiore, callo tenui, subdiffuso obducto; apertura diagonali, triangulari-semiovali; perist. albo, breviter expanso, margine basali leviter arcuato, angulatim cum columella brevi, simplice juncto.* —Operc. ?

Diam. maj. $5\frac{1}{2}$, min. $4\frac{1}{2}$, alt. $3\frac{1}{2}$ mill.

Hab. Drayton Range, North Australia.

32. *HELICINA HEATEI*, Pfr. *H. testa subdepressa, tenuiuscula, striatula et subgranulata, parum nitida, fasciis latis saturatae purpureis et albis picta; spira brevissime conoidea, mucronulata; anfr. 5 vix convexiusculis, ultimo lato, peripheria subcarinato, basi circa callum fulvo-aurantiacum, circumscriptum albo; apertura perobliqua, triangulari-semiovali; columella brevi, perarcuata; perist. reflexo, late aurantiaco, margine basali levissime arcuato, immediate in callum basalem continuato.* —Operc. ?

Diam. maj. 14, min. $11\frac{1}{3}$, alt. $7\frac{1}{2}$ mill.

β. *Lutea, spira et fascia unica anfr. ultimi supera purpureis, callo basali et peristomate igneis.*

Hab. Island of Granada, West Indies (named after R. W. Heate, Esq., Lieut.-Governor of the Island).

33. *HELICINA RUFa*, Pfr. *H. testa subconoideo-depressa, solidula, subrugoso-striata et punctulata, nitida, rufa; spira subconoideo-convexa, vertice subtili; anfr. $4\frac{1}{2}$, superis planiusculis, ultimo lato, depresso, peripheria subrotundato; apertura diagonali, late semiovali; columella verticaliter ab anfr. penultimo descendente, leviter curvata, antice in tuberculum desinente, callum basalem emitente subgranulatum, circumscriptum; perist. breviter expanso, albo-limbato.* —Operc. concolor.

Diam. maj. 13, min. 11, alt. $6\frac{2}{3}$ mill.

β. *Pallide straminea, perist. intus pallide aurantiaco.*

γ. *Minor, rufa.*

Diam. maj. 10, min. $8\frac{2}{3}$, alt. $5\frac{1}{2}$ mill.

Hab. Haiti (Mr. Salle).